



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

and, consequently, the $\triangle MNO = \frac{1}{8}(\mathbf{N})^2 / \sin A \sin B \sin C \dots (5)$. The expression for the average area of the $\triangle MNO$, therefore, becomes

$$\mathbf{A} = \frac{1}{8abc \sin A \sin B \sin C} \int_0^a \int_0^b \int_0^c (\mathbf{N})^2 dx dy dz$$

$$= \frac{1}{24} \left(\frac{a^2 \sin A}{\sin B \sin C} + \frac{b^2 \sin B}{\sin C \sin A} + \frac{c^2 \sin C}{\sin A \sin B} \right) = \frac{a^4 + b^4 + c^4}{48 \Delta}.$$

[Note.—Problems twenty-three and twenty-four are identical. This fact was not observed until after they were both printed. ED.]

PROBLEMS.

31. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

Find the average length of a line drawn at random across the opposite sides of a rectangle whose length is l and breadth b .

32. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

Find the average area of the random sector whose vertex is a random point in a given circle.

MISCELLANEOUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

17. Proposed by SAMUEL HART WRIGHT, M. D., M. A., Ph. D. Penn Yan, New York.

A bright star passed my meridian at 7 P. M. The Chronometer soon after ran down and stopped, but I set it again when the same star had a true altitude of $30^\circ = \alpha$. What time was it then, my latitude being $42^\circ 30' \text{ N.} = \lambda$, and the star's declination $60^\circ \text{ N.} = \delta$

II. Solution by the PROPOSER.

Let B be the north pole, A the zenith, C the star, HH' the horizon, AH and AH' each $= 90^\circ$, AH' being a meridian, AH a verticle circle, BH' the altitude of the pole = the latitude $= L$, $AB = \text{co-latitude} = c$, $BC = a = \text{polar distance of the star} = P$, $AC = b = \text{the zenith distance of star}$, $CH = A = \text{altitude of the star}$, and the angle $ABC = \text{the hour-angle of the star} = T$ in sidereal time. Put $s = \frac{1}{2}(a + b + c)$, and $s - a = a'$, $s - b = b'$, and $s - c = c'$. Then by *Sph. Trig.*

$$\sin \frac{1}{2} T = \sqrt{\left(\frac{\sin c' \sin a'}{\sin c \sin a} \right)}, \text{ and}$$



